

DR CHARLIE BOAST BDS(Adel), D.Clin.Dent (Qld), FRACDS

REGISTERED DENTAL SPECIALISTS

TWO-IMPLANT LOWER OVERDENTURE

Simple to make, Easy to wear.



"The 1st choice standard of care for edentulous patients"

The Magill consensus statement on overdentures: Int J Oral Maxfac 2002; Int J Prosthetics 2002; Gerodontology 2002

Dental implants are titanium screws that when placed appropriately in the jaw become fused (osseo-integrated) with the living bone and can retain and stabilise dentures or even support fixed crowns and bridgework. There is over 40 years of solid medical data recording their long-term success in dentistry.

Patients who have lost all their teeth (edentulous) are a particular needy group as their lower full dentures are renowned for being unstable and difficult to wear, which may then compromise good nutrition and quality of life. The two implant-retained mandibular over-denture significantly increases chewing function (approximately sixfold) and has been considered the "Ist choice standard of care for lower edentulous patients" for some time now (The Magill Consensus Statement on Overdentures: *Int J Oral Maxfac 2002; Int J Prosthetics 2002; Gerodontology 2002)*.

PROTOCOL

(Periodontists/ Surgical Total fee \$5561 + Prosthesist fee)

- * Bookings / Enquiries: Pacific Periodontics will post an information pack out.
- 1) Patient brings old dentures and toothbrush to consultation appointment.

Item no. 015

Item no. 039 (3-D conebeam scan)

Appointment involves:

- Assessment of medical history and oral/gum health
- Quality, quantity and shape of residual jaw bone and relationship of Mandibular nerve assessed clinically and radiographically (via Cone Beam scan).
- Explanation of risks and benefit of surgical procedure.
- Pre-operative medications (if needed) dispensed.

Two Implant Lower Overdenture

Cost \$104

Cost \$157



DR CHARLIE BOAST BDS(Adel), D.Clin.Dent (Qld), FRACDS

REGISTERED DENTAL SPECIALISTS

2) Surgical placement of just two titanium implants and using local anaesthetic +/- oral Lorazepam sedation. Implants are kept submerged under gum to allow continued wear of dentures.
Item no.:- 684 x 2, 022 x 2, 927 Cost \$3800
3) 3 months later:- Minor surgical exposure of implants and placement of healing abutment.
Item no:- 012, 022 x 2, 661 x 2 Cost \$750

4) 1 week later:- Place gold equator locator abutments.

ltem no:- 691 x 2

- 5) On same day Patient returns and pays Prosthetist to:
- · Convert existing denture into an "implant-retained overdenture". Or
- Make new lower "implant-retained overdenture"

MAINTENANCE

"Significantly fewer visits for adjustment related to post-placement pressure spots were required for mandibular overdentures than for conventional mandibular full dentures."

Klemetti E et al. J Can Dent Assoc 2003; 69(1):29-33

Similar to conventional full dentures, an implant retained over-denture will require occasional maintenance. The Equator/Locator caps have been tested to 56,000 cycles in the laboratory, however may require adjustment or replacement after a couple of years by the prosthetist.

The dental implants should be checked clinically by a dentist on a yearly basis and an x-ray taken 12 months after initial use, then on a 3 yearly basis.

Why choose Pacific Periodontics and Implants?

- Board Registered Specialists
- In house 3D Scan for digital virtual guidance ensuring accurate and safe placement
- We only use 100% commercially pure titanium quality implants
- I0 year warranty

Two Implant Lower Overdenture





Cost \$750